


# Climate4you update October 2018




## Contents:

- Page 2: October 2018 global surface air temperature overview
- Page 3: Comments to the October 2018 global surface air temperature overview
- Page 4: Temperature quality class 1: Lower troposphere temperature from satellites
- Page 5: Temperature quality class 2: HadCRUT global surface air temperature
- Page 6: Temperature quality class 3: GISS and NCDC global surface air temperature
- Page 9: Comparing global surface air temperature and satellite-based temperatures
- Page 10: Global air temperature linear trends
- Page 11: Global temperatures: All in one, Quality Class 1, 2 and 3
- Page 13: Global sea surface temperature
- Page 16: Ocean temperature in uppermost 100 m
- Page 18: North Atlantic heat content uppermost 700 m
- Page 19: North Atlantic temperatures 0-800 m depth along 59N, 30-0W
- Page 20: Global ocean temperature 0-1900 m depth summary
- Page 21: Global ocean net temperature change since 2004 at different depths
- Page 22: La Niña and El Niño episodes
- Page 23: Troposphere and stratosphere temperatures from satellites
- Page 24: Zonal lower troposphere temperatures from satellites
- Page 25: Arctic and Antarctic lower troposphere temperatures from satellites
- Page 26: Temperature over land versus over oceans
- Page 27: Arctic and Antarctic surface air temperatures
- Page 30: Arctic and Antarctic sea ice
- Page 34: Sea level in general
- Page 35: Global sea level from satellite altimetry
- Page 36: Global sea level from tide gauges
- Page 37: Northern Hemisphere weekly and seasonal snow cover
- Page 39: Atmospheric specific humidity
- Page 40: Atmospheric CO<sub>2</sub>
- Page 41: Relation between annual change of atm. CO<sub>2</sub> and La Niña and El Niño episodes
- Page 42: Phase relation between atmospheric CO<sub>2</sub> and global temperature
- Page 43: Global air temperature and atmospheric CO<sub>2</sub>
- Page 47: Latest 20-year QC1 global monthly air temperature change
- Page 48: Sunspot activity and QC1 average satellite global air temperature
- Page 49: Climate and history: 2018. *The concept of climate revisited.*

## October 2018 global surface air temperature overview



2



October 2018 surface air temperature compared to the average of the last 10 years. Green-yellow-red colours indicate areas with higher temperature than the 10-year average, while blue colours indicate lower than average temperatures. Data source: [Goddard Institute for Space Studies](#) (GISS) using Hadl\_Reyn\_v2 ocean surface temperatures.

## Comments to the October 2018 global surface air temperature overview

General: This newsletter contains graphs showing a selection of key meteorological variables for the past month. All temperatures are given in degrees Celsius.

In the above maps showing the geographical pattern of surface air temperatures, the last previous 10 years are used as reference period.

The rationale for comparing with this recent period instead of the official WMO 'normal' period 1961-1990, is that the latter period is affected by the cold period 1945-1980. Most comparisons with this time period will automatically appear as warm, and it will be difficult to decide if modern surface air temperatures are increasing or decreasing. Comparing instead with the last previous 10 years overcomes this problem and displays the modern dynamics of ongoing change. This decadal approach also corresponds well to the typical memory horizon for many people.

3

In addition, the GISS temperature data used for preparing the above diagrams display distinct temporal instability for data before the turn of the century (see p. 7). Any comparison with the WMO 'normal' period 1961-1990 is therefore influenced by ongoing monthly mainly administrative changes. An unstable value is clearly not suited as reference value. Simply comparing with the last previous 10 years makes more sense and is more useful.

The different air temperature records have been divided into three quality classes, QC1, QC2 and QC3, respectively, as described on page 7.

In many diagrams shown in this newsletter the thin line represents the monthly global average value, and the thick line indicate a simple running average, in most cases a simple moving 37-month average, nearly corresponding to a three-year average. The 37-month average is calculated from values covering a range from 18 months before to 18 months after, with equal weight given to all individual months.

The year 1979 has been chosen as starting point in many diagrams, as this roughly corresponds to both the beginning of satellite observations and the onset of the late 20<sup>th</sup> century warming period. However, several of the data series have a much longer record length, which may be inspected in greater detail on [www.climate4you.com](http://www.climate4you.com).

### October 2018 global surface air temperatures


General: For October 2018 GISS supplied 15931 interpolated surface air data points; all values are used to produce the diagrams shown on page 2. According to the GISS data, the average global September temperature anomaly was higher than in the previous months. However, the overall decrease of global surface air temperatures since the 2016-17 El Niño global temperature peak persists, as shown by the diagrams on p. 4-6.

The Northern Hemisphere anomaly pattern was characterised by large regional contrasts. Most of Northern America and Greenland were cold compared to the previous 10 years, with only Alaska being relatively warm. In contrast to North America, most of Europe, Russia (incl. Siberia) was relatively warm. The peak warm anomaly was in northern Siberia. The Canadian-Greenland sector of the Arctic was cold, while the Europe-Siberia-Alaska sector was warm. However, the GISS surface air temperatures north of 80° N still appears to have an issue with interpolation, resulting in a unnatural circular pattern.

Near the Equator temperatures were near or below the 10-year average.


The Southern Hemisphere temperatures were near the average for the previous 10 years. Only South Africa and northern Australia was relatively warm. South America was mainly below the 10-yr average. In the Antarctica, most regions were relatively warm, although the entire Atlantic sector was relatively cold. South of 80°S the GISS anomaly pattern is still influenced by an interpolation artefact.

## Temperature quality class 1: Lower troposphere temperature from satellites, updated to October 2018




Global monthly average lower troposphere temperature (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37-month average.

4



Global monthly average lower troposphere temperature (thin line) since 1979 according to according to [Remote Sensing Systems](#) (RSS), USA. The thick line is the simple running 37-month average.


## Temperature quality class 2: HadCRUT global surface air temperature, updated to September 2018



5


Global monthly average surface air temperature (thin line) since 1979 according to according to the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK. The thick line is the simple running 37-month average.

### Temperature quality class 3: GISS and NCDC global surface air temperature, updated to October 2018



Global monthly average surface air temperature (thin line) since 1979 according to according to the [Goddard Institute for Space Studies \(GISS\)](#), at Columbia University, New York City, USA, using ERSST\_v4 ocean surface temperatures. The thick line is the simple running 37-month average.

6



Global monthly average surface air temperature since 1979 according to according to the [National Climatic Data Center \(NCDC\)](#), USA. The thick line is the simple running 37-month average.

### A note on data record stability and -quality:

All temperature diagrams shown above have 1979 as starting year. This roughly marks the beginning of the recent period of global warming, after termination of the previous period of global cooling from about 1940. In addition, the year 1979 also represents the starting date for the satellite-based global temperature estimates (UAH and RSS). For the three surface air temperature records (HadCRUT, NCDC and GISS), they begin much earlier (in 1850 and 1880, respectively), as can be inspected on [www.climate4you.com](http://www.climate4you.com).

For all three surface air temperature records, but especially NCDC and GISS, administrative changes to anomaly values are quite often introduced, even for observations many years back in time. Some changes may be due to the delayed addition of new station data or change of station location, while others probably have their origin in a change of technique to calculate average values. It is clearly impossible to evaluate the validity of such administrative changes for the outside user of these records; it is only possible to note that such changes appear very often (see example diagram next page).

In addition, the three surface records represent a blend of sea surface data collected by moving ships or by other means, plus data from land stations of partly unknown quality and unknown degree of representativeness for their region. Many of the land stations have also moved geographically during their existence, and their instrumentation changed, and they are influenced by changes in their surroundings (vegetation, buildings, etc.).

The satellite temperature records also have their problems, but these are generally of a more technical nature and therefore correctable. In addition, the temperature sampling by satellites is more regular and complete on a global basis than that represented by the surface records. Also important is that the sensors on satellites measure

temperature directly by emitted radiation, while most surface temperature measurements are indirect, using electronic resistance.

Everybody interested in climate science should gratefully acknowledge the efforts put into maintaining the different temperature databases referred to in the present newsletter. At the same time, however, it is also important to realise that all temperature records cannot be of equal scientific quality. The simple fact that they to some degree differ shows that they cannot all be correct.

On this background, and for practical reasons, Climate4you operates with three quality classes (1-3) for global temperature records, with 1 representing the highest quality level:

Quality class 1: The satellite records (UAH and RSS).

Quality class 2: The HadCRUT surface record.

Quality class 3: The NCDC and GISS surface records.

The main reason for discriminating between the three surface records is the following:

While both NCDC and GISS often experience quite large administrative changes (see example on p.8), and therefore essentially are unstable temperature records, the changes introduced to HadCRUT are fewer and smaller. For obvious reasons, as the past does not change, any record undergoing continuing changes cannot describe the past correctly all the time. Frequent and large corrections in a database also signal fundamental doubt about what is likely to represent the correct values.

You can find more on the issue of lack of temporal stability on [www.climate4you.com](http://www.climate4you.com) (go to: *Global Temperature*, followed by *Temporal Stability*).





Diagram showing the adjustment made since May 2008 by the [Goddard Institute for Space Studies](#) (GISS), USA, in anomaly values for the months January 1910 and January 2000.

**Note:** The administrative upsurge of the temperature increase from January 1915 to January 2000 has grown from 0.45 (reported May 2008) to 0.67°C (reported November 2018). This represents an about 49% administrative temperature increase over this period, meaning that about half of the apparent (as reported by GISS) global temperature increase from January 1910 to January 2000 is due to administrative changes of the original data since May 2008.

**Comparing global surface air temperature and lower troposphere satellite temperatures; updated to September 2018**



9

*Plot showing the average of monthly global surface air temperature estimates ([HadCRUT4](#), [GISS](#) and [NCDC](#)) and satellite-based temperature estimates ([RSS MSU](#) and [UAH MSU](#)). The thin lines indicate the monthly value, while the thick lines represent the simple running 37-month average, nearly corresponding to a running 3-yr average. The lower panel shows the monthly difference between average surface air temperature and satellite temperatures. As the base period differs for the different temperature estimates, they have all been normalised by comparing to the average value of 30 years from January 1979 to December 2008.*

NOTE: Since about 2003, the average global surface air temperature has been drifting away in positive direction from the average satellite temperature, meaning that the surface records show warming in relation to the troposphere records. The reason(s) for this is not entirely clear but can presumably at least partly be explained by the recurrent administrative adjustments made to the surface records (see p. 7-8).

## Global air temperature linear trends updated to September 2018




Diagram showing the latest 5, 10, 20 and 30-yr linear annual global temperature trend, calculated as the slope of the linear regression line through the data points, for two satellite-based temperature estimates (UAH MSU and RSS MSU).

10






Diagram showing the latest 5, 10, 20, 30, 50, 70 and 100-year linear annual global temperature trend, calculated as the slope of the linear regression line through the data points, for three surface-based temperature estimates (GISS, NCDC and HadCRUT4).

**All in one, Quality Class 1, 2 and 3; updated to September 2018**




Superimposed plot of Quality Class 1 (UAH and RSS) global monthly temperature estimates. As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of both temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.

11



Superimposed plot of Quality Class 1 and 2 (UAH, RSS and HadCRUT4) global monthly temperature estimates. As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of all three temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.



*Superimposed plot of Quality Class 1, 2 and 3 global monthly temperature estimates (UAH, RSS, HadCRUT4, GISS and NCDP). As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of all five temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.*

Please see notes on page 7 relating to the above three quality classes.


It should be kept in mind that satellite- and surface-based temperature estimates are derived from different types of measurements, and that comparing them directly as done in the diagram above therefore may be somewhat problematical. However, as both types of estimate often are discussed together, the above diagram may nevertheless be of some interest. In fact, the different types of temperature estimates appear to agree as to the overall temperature variations on a 2-3-year scale, although on a shorter time scale there are often considerable differences between the individual records. However, since about 2003 the surface records are slowly drifting towards higher temperatures than the combined satellite record (see p. 9).

The average of all five global temperature estimates presently shows an overall stagnation, at least since

2002-2003. There has been no significant increase in global air temperature since 1998, which however was affected by the oceanographic El Niño event. Also, the recent (2015-16) El Niño event may well represent a relatively short-lived spike on a longer development. The coming years will show if this is the case or not. Neither has there been a temperature decrease since about 2002-2003.

The present temperature stagnation does not exclude the possibility that global temperatures will begin to increase significantly again later. On the other hand, it also remains a possibility that Earth just now is passing an overall temperature peak, and that global temperatures will begin to decrease during the coming years. Again, time will show which of these possibilities is correct.

## Global sea surface temperature, updated to October 2018



13

*Sea surface temperature anomaly on 26 October 2018. Map source: Plymouth State Weather Center. Reference period: 1977-1991.*


Because of the large surface areas near Equator, the temperature of the surface water in these regions is especially important for the global atmospheric temperature (p. 4-6). No less than 50% of planet Earth's surface area is located within 30°N and 30°S.

A mixture of relatively warm and cold water in August 2018 dominates much of the oceans near the Equator. In addition, parts of the North Atlantic are cooling. Presumably, all this will be influencing global air temperatures in the months to come.

However, the significance of any short-term cooling or warming reflected in air temperatures should not


be overstated. Whenever Earth experiences cold La Niña or warm El Niño episodes (Pacific Ocean) major heat exchanges takes place between the Pacific Ocean and the atmosphere above, eventually showing up in estimates of the global air temperature.

However, this does not reflect similar changes in the total heat content of the atmosphere-ocean system. In fact, global net changes can be small and such heat exchanges may mainly reflect redistribution of energy between ocean and atmosphere. What matters is the overall temperature development when seen over several years.




Global monthly average lower troposphere temperature over oceans (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats.

14




Global monthly average sea surface temperature since 1979 according to University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK. Base period: 1961-1990. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats.



Global monthly average sea surface temperature since 1979 according to the [National Climatic Data Center](#) (NCDC), USA. Base period: 1901-2000. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats.


**June 18, 2015:** NCDC has introduced several rather large administrative changes to their sea surface temperature record. The overall result is to produce a record giving the impression of a continuous temperature increase, also in the 21st century. As the oceans cover about 71% of the entire surface of planet Earth, the effect of this administrative change is clearly seen in the NCDC record for global surface air temperature (p. 6).

## Ocean temperature in uppermost 100 m, updated to September 2018




World Oceans vertical average temperature 0-100 m depth since 1955. The thin line indicates 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.


16



Pacific Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicate 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.




Atlantic Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicate 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](https://www.noaa.gov/data/ocean/obs/sea-surface/sea-surface-temperature-anomaly). Base period 1955-2010.




Indian Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicate 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](https://www.noaa.gov/data/ocean/obs/sea-surface/sea-surface-temperature-anomaly). Base period 1955-2010.

## North Atlantic heat content uppermost 700 m, updated to September 2018




18




Global monthly heat content anomaly ( $10^{18}$  Joules) in the uppermost 700 m of the North Atlantic (60-0W, 30-65N; see map above) ocean since January 1955. The thin line indicates monthly values, and the thick line represents the simple running 37-month (c. 3 year) average. Data source: [National Oceanographic Data Center](#) (NODC).

**North Atlantic temperatures 0-800 m depth along 59°N, 30-0W, updated to June 2018**




Time series depth-temperature diagram along 59 N across the North Atlantic Current from 30°W to 0°W, from surface to 800 m depth. Source: [Global Marine Argo Atlas](#). See also the diagram below.

19




Average temperature along 59 N, 30-0W, 0-800m depth, corresponding to the main part of the North Atlantic Current, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100.

## Global ocean temperature 0-1900 m depth summary, updated to June 2018




Summary of average temperature in uppermost 1900 m in different parts of the global oceans, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100.

Global ocean net temperature change since 2004 at different depths, updated to June 2018




Net temperature change since 2004 from surface to 1900 m depth in different parts of the global oceans, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100. Please note that due to the spherical form of Earth, northern and southern latitudes represent only small ocean volumes, compared to latitudes near the Equator.

## La Niña and El Niño episodes, updated to October 2018




Warm ( $>+0.5^{\circ}\text{C}$ ) and cold ( $<-0.5^{\circ}\text{C}$ ) episodes for the [Oceanic Niño Index \(ONI\)](#), defined as 3 month running mean of ERSSTv4 SST anomalies in the Niño 3.4 region ( $5^{\circ}\text{N}$ - $5^{\circ}\text{S}$ ,  $120^{\circ}$ - $170^{\circ}\text{W}$ ). For historical purposes cold and warm episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons. Anomalies are centred on 30-yr base periods updated every 5 years.

## Troposphere and stratosphere temperatures from satellites, updated to October 2018




Global monthly average temperature in different according to University of Alabama at Huntsville, USA. The thin lines represent the monthly average, and the thick line the simple running 37-month average, nearly corresponding to a running 3-year average.

## Zonal lower troposphere temperatures from satellites, updated to October 2018




Global monthly average lower troposphere temperature since 1979 for the tropics and the northern and southern extratropics, according to University of Alabama at Huntsville, USA. Thin lines show the monthly temperature. Thick lines represent the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.

## Arctic and Antarctic lower troposphere temperature, updated to October 2018



Global monthly average lower troposphere temperature since 1979 for the North Pole and South Pole regions, based on satellite observations ([University of Alabama](#) at Huntsville, USA). Thin lines show the monthly temperature. The thick line is the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.

## Temperature over land versus over oceans, updated to October 2018



Global monthly average lower troposphere temperature since 1979 measured over land and oceans, respectively, according to [University of Alabama](#) at Huntsville, USA. Thick lines are the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.

## Arctic and Antarctic surface air temperature, updated to September 2018




Diagram showing area weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 2000, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.

27




Diagram showing area weighted Antarctic (70-90°S) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 2000, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1957, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area weighted Antarctic (70-90°S) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1957, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area-weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1920, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.

Because of the relatively small number of Arctic stations before 1930, month-to-month variations in the early part of the Arctic temperature record 1920-2018 are larger than later (diagram above).

The period from about 1930 saw the establishment of many new Arctic meteorological stations, first in Russia and Siberia, and following the 2nd World War, also in North America. The period since 2005 is warm, about as warm as the period 1930-1940.


As the HadCRUT4 data series has improved high latitude coverage data coverage (compared to the HadCRUT3 series), the individual 5°x5° grid cells has been weighted according to their surface area. This

contrasts with [Gillett et al. 2008](#) which calculated a simple average, with no correction for the significant surface area effect of latitude in polar regions.

Literature:


Gillett, N.P., Stone, D.A., Stott, P.A., Nozawa, T., Karpechko, A.Y.U., Hegerl, G.C., Wehner, M.F. and Jones, P.D. 2008. Attribution of polar warming to human influence. *Nature Geoscience* 1, 750-754.

## Arctic and Antarctic sea ice, updated to October 2018




30

Sea ice extent 26 October 2018. The median limit of sea ice (orange line) is defined as 15% sea ice cover, according to the average of satellite observations 1981-2010 (both years included). Sea ice may therefore well be encountered outside and open water areas inside the limit shown in the diagrams above. Map source: National Snow and Ice Data Center (NSIDC).



Diagrams showing Arctic sea ice extent and concentration 27 October 2017 (left) and 2018 (right), according to the Japan Aerospace Exploration Agency (JAXA).



Graphs showing monthly Antarctic, Arctic and global sea ice extent since November 1978, according to the [National Snow and Ice data Center \(NSIDC\)](#).

31






Diagram showing daily Arctic sea ice extent since June 2002, to 27 October 2018, by courtesy of [Japan Aerospace Exploration Agency \(JAXA\)](#).


## Sea Ice Thickness, 27-Oct-2018



32



Diagrams showing Arctic sea ice extent and thickness 27 October 2017 (left) and 2018 (right and above) and the seasonal cycles of the calculated total arctic sea ice volume, according to [The Danish Meteorological Institute \(DMI\)](#). The mean sea ice volume and standard deviation for the period 2004-2013 are shown by grey shading.



12 month running average sea ice extension, global and in both hemispheres since 1979, the satellite-era. The October 1979 value represents the monthly 12-month average of November 1978 - October 1979, the November 1979 value represents the average of December 1978 - November 1979, etc. The stippled lines represent a 61-month (ca. 5 years) average. Data source: National Snow and Ice Data Center (NSIDC).

## Sea level in general

Global (or eustatic) sea-level change is measured relative to an idealised reference level, the geoid, which is a mathematical model of planet Earth's surface (Carter et al. 2014). Global sea-level is a function of the volume of the ocean basins and the volume of water they contain. Changes in global sea-level are caused by – but not limited to - four main mechanisms:

1. Changes in local and regional air pressure and wind, and tidal changes introduced by the Moon.
2. Changes in ocean basin volume by tectonic (geological) forces.
3. Changes in ocean water density caused by variations in currents, water temperature and salinity.
4. Changes in the volume of water caused by changes in the mass balance of terrestrial glaciers.

In addition to these there are other mechanisms influencing sea-level; such as storage of ground water, storage in lakes and rivers, evaporation, etc.

Mechanism 1 is controlling sea-level at many sites on a time scale from months to several years. As an example, many coastal stations show a pronounced annual variation reflecting seasonal changes in air pressures and wind speed. Longer-term climatic changes playing out over decades or centuries will also affect measurements of sea-level changes. Hansen et al. (2011, 2015) provide excellent analyses of sea-level changes caused by recurrent changes of the orbit of the Moon and other phenomena.

Mechanism 2 – with the important exception of earthquakes and tsunamis - typically operates over long (geological) time scales, and is not significant on human time scales. It may relate to variations in the sea-floor spreading rate, causing volume changes in mid-ocean mountain ridges, and to the slowly changing configuration of land and oceans. Another effect may be the slow rise of basins due to isostatic offloading by deglaciation after an ice age. The floor of the Baltic Sea and the Hudson Bay are presently rising, causing a slow net transfer of

water from these basins into the adjoining oceans. Slow changes of very big glaciers (ice sheets) and movements in the mantle will affect the gravity field and thereby the vertical position of the ocean surface. Any increase of the total water mass as well as sediment deposition into oceans increase the load on their bottom, generating sinking by viscoelastic flow in the mantle below. The mantle flow is directed towards the surrounding land areas, which will rise, thereby partly compensating for the initial sea level increase induced by the increased water mass in the ocean.

Mechanism 3 (temperature-driven expansion) only affects the uppermost part of the oceans on human time scales. Usually, temperature-driven changes in density are more important than salinity-driven changes. Seawater is characterised by a relatively small coefficient of expansion, but the effect should however not be overlooked, especially when interpreting satellite altimetry data. Temperature-driven expansion of a column of seawater will not affect the total mass of water within the column considered and will therefore not affect the potential at the top of the water column. Temperature-driven ocean water expansion will therefore not in itself lead to lateral displacement of water, but only lift the ocean surface locally. Near the coast, where people are living, the depth of water approaches zero, so no temperature-driven expansion will take place here (Mörner 2015). Mechanism 3 is for that reason not important for coastal regions.


Mechanism 4 (changes in glacier mass balance) is an important driver for global sea-level changes along coasts, for human time scales. Volume changes of floating glaciers – ice shelves – has no influence on the global sea-level, just like volume changes of floating sea ice has no influence. Only the mass-balance of grounded or land-based glaciers is important for the global sea-level along coasts.

Summing up: Mechanism 1 and 4 are the most important for understanding sea-level changes along coasts.

### References:


- Carter R.M., de Lange W., Hansen, J.M., Humlum O., Idso C., Kear, D., Legates, D., Mörner, N.A., Ollier C., Singer F. & Soon W. 2014. Commentary and Analysis on the Whitehead & Associates 2014 NSW Sea-Level Report. Policy Brief, NIPCC, 24. September 2014, 44 pp. <http://climatechangereconsidered.org/wp-content/uploads/2014/09/NIPCC-Report-on-NSW-Coastal-SL-9z-corrected.pdf>
- Hansen, J.-M., Aagaard, T. and Binderup, M. 2011. Absolute sea levels and isostatic changes of the eastern North Sea to central Baltic region during the last 900 years. *Boreas*, 10.1111/j.1502-3885.2011.00229.x. ISSN 0300-9483.
- Hansen, J.-M., Aagaard, T. and Huijpers, A. 2015. Sea-Level Forcing by Synchronization of 56- and 74-Year Oscillations with the Moon's Nodal Tide on the Northwest European Shelf (Eastern North Sea to Central Baltic Sea). *Journ. Coastal Research*, 16 pp.
- Mörner, Nils-Axel 2015. Sea Level Changes as recorded in nature itself. *Journal of Engineering Research and Applications*, Vol.5, 1, 124-129.

## Global sea level from satellite altimetry, updated to January 2018



Global sea level since December 1992 according to the Colorado Center for Astrodynamics Research at University of Colorado at Boulder. The blue dots are the individual observations, and the purple line represents the running 121-month (ca. 10 year) average. The two lower panels show the annual sea level change, calculated for 1 and 10-year time windows, respectively. These values are plotted at the end of the interval considered. Data from the TOPEX/Poseidon mission have been used before 2002, and data from the Jason-1 mission (satellite launched December 2001) after 2002.

## Global sea level from tide-gauges, updated to December 2017




36

*Holgate-9* monthly tide gauge data from PSMSL Data Explorer. *Holgate (2007)* suggested the nine stations listed in the diagram to capture the variability found in a larger number of stations over the last half century studied previously. For that reason, average values of the *Holgate-9* group of tide gauge stations are interesting to follow. The blue dots are the individual average monthly observations, and the purple line represents the running 121-month (ca. 10 year) average. The two lower panels show the annual sea level change, calculated for 1 and 10-year windows, respectively. These values are plotted at the end of the interval considered.

### Reference:


Holgate, S.J. 2007. On the decadal rates of sea level change during the twentieth century. *Geophys. Res. Letters*, 34, L01602, doi:10.1029/2006GL028492

**Northern Hemisphere weekly and seasonal snow cover, updated to October 2018**




Northern hemisphere snow cover (white) and sea ice (yellow) 27 October 2017 (left) and 2018 (right). Map source: [National Ice Center \(NIC\)](#).


37



Northern hemisphere weekly snow cover since January 2000 according to Rutgers University Global Snow Laboratory. The thin blue line is the weekly data, and the thick blue line is the running 53-week average (approximately 1 year). The horizontal red line is the 1972-2017 average.




Northern hemisphere weekly snow cover since January 1972 according to Rutgers University Global Snow Laboratory. The thin blue line is the weekly data, and the thick blue line is the running 53-week average (approximately 1 year). The horizontal red line is the 1972-2017 average.




Northern hemisphere seasonal snow cover since January 1972 according to Rutgers University Global Snow Laboratory.

## Atmospheric specific humidity, updated to October 2018




*Specific atmospheric humidity (g/kg) at three different altitudes in the lower part of the atmosphere (the Troposphere) since January 1948 (Kalnay et al. 1996). The thin blue lines show monthly values, while the thick blue lines show the running 37-month average (about 3 years). Data source: [Earth System Research Laboratory \(NOAA\)](#).*

## Atmospheric CO<sub>2</sub>, updated to October 2018




Monthly amount of atmospheric CO<sub>2</sub> (upper diagram) and annual growth rate (lower diagram); average last 12 months minus average preceding 12 months, thin line) of atmospheric CO<sub>2</sub> since 1959, according to data provided by the [Mauna Loa Observatory](#), Hawaii, USA. The thick, stippled line is the simple running 37-observation average, nearly corresponding to a running 3-year average.

## The relation between annual change of atmospheric CO<sub>2</sub> and La Niña and El Niño episodes, updated to October 2018



*Visual association between annual growth rate of atmospheric CO<sub>2</sub> (upper panel) and Oceanic Niño Index (lower panel). See also diagrams on page 40 and 22, respectively.*

## The phase relation between atmospheric CO<sub>2</sub> and global temperature, updated to September 2018




12-month change of global atmospheric CO<sub>2</sub> concentration ([Mauna Loa](#); green), global sea surface temperature ([HadSST3](#); blue) and global surface air temperature ([HadCRUT4](#); red dotted). All graphs are showing monthly values of DIFF12, the difference between the average of the last 12 month and the average for the previous 12 months for each data series.


### References:



Humlum, O., Stordahl, K. and Solheim, J-E. 2012. The phase relation between atmospheric carbon dioxide and global temperature. *Global and Planetary Change*, August 30, 2012.  
<http://www.sciencedirect.com/science/article/pii/S0921818112001658?v=s5>


# Global air temperature and atmospheric CO<sub>2</sub>, updated to October 2018



43







Diagrams showing UAH, RSS, HadCRUT4, GISS, and NDCC monthly global air temperature estimates (blue) and the monthly atmospheric CO<sub>2</sub> content (red) according to the [Mauna Loa Observatory](#), Hawaii. The Mauna Loa data series begins in March 1958, and 1958 was therefore chosen as starting year for the diagrams. Reconstructions of past atmospheric CO<sub>2</sub> concentrations (before 1958) are not incorporated in this diagram, as such past CO<sub>2</sub> values are derived by other means (ice cores, stomata, or older measurements using different methodology), and therefore are not directly comparable with direct atmospheric measurements. The dotted grey line indicates the approximate linear temperature trend, and the boxes in the lower part of the diagram indicate the relation between atmospheric CO<sub>2</sub> and global surface air temperature, negative or positive. Please note that the HadCRUT diagram (p.43) is not yet updated beyond September 2018.

Most climate models are programmed to give the greenhouse gas carbon dioxide CO<sub>2</sub> important influence on global temperature. It is therefore relevant to compare different temperature records with measurements of atmospheric CO<sub>2</sub>, as shown in the diagrams above.

Any comparison, however, should not be made on a monthly or annual basis, but for a longer time, as other effects (oceanographic, etc.) may well override the potential influence of CO<sub>2</sub> on short time scales such as just a few years. It is of cause equally inappropriate to present new meteorological record values, whether daily, monthly or annual, as support for the hypothesis ascribing high importance of atmospheric CO<sub>2</sub> for global temperatures. Any such

meteorological record value may well be the result of other phenomena.

What exactly defines the critical length of a relevant period length to consider for evaluating the alleged importance of CO<sub>2</sub> remains elusive and represents a theme for discussion. However, the length of the critical period must be inversely proportional to the temperature sensitivity of CO<sub>2</sub>, including feedback effects. If the net temperature effect of atmospheric CO<sub>2</sub> is strong, the critical period will be short, and vice versa.

However, past climate research history provides some clues as to what has traditionally been considered the relevant length of period over which


to compare temperature and atmospheric CO<sub>2</sub>. After about 10 years of concurrent global temperature- and CO<sub>2</sub>-increase, IPCC was established in 1988. For obtaining public and political support for the CO<sub>2</sub>-hypothesis the 10-year warming period leading up to 1988 most likely was important. Had the global temperature instead been decreasing, political support for the hypothesis would have been difficult to obtain.

Based on the previous 10 years of concurrent temperature- and CO<sub>2</sub>-increase, many climate scientists in 1988 presumably felt that their

understanding of climate dynamics was sufficient to conclude about the importance of CO<sub>2</sub> for global temperature changes. From this it may safely be concluded that 10 years was considered a period long enough to demonstrate the effect of increasing atmospheric CO<sub>2</sub> on global temperatures.

Adopting this approach as to critical time length (at least 10 years), the varying relation (positive or negative) between global temperature and atmospheric CO<sub>2</sub> has been indicated in the lower panels of the diagrams above.

## Latest 20-year QC1 global monthly air temperature changes, updated to October 2018



47

Last 20 years' global monthly average air temperature according to Quality Class 1 (UAH and RSS; see p.10) global monthly temperature estimates. The thin blue line represents the monthly values. The thick black line is the linear fit, with 95% confidence intervals indicated by the two thin black lines. The thick green line represents a 5-degree polynomial fit, with 95% confidence intervals indicated by the two thin green lines. A few key statistics are given in the lower part of the diagram (please note that the linear trend is the monthly trend).


In the ongoing climate debate the following about the global surface air temperature is often put forward: Is the surface air temperature still increasing or has it basically remained without significant changes during the last about 15 years?

The diagram above may be useful in this context and demonstrates the differences between two often used statistical approaches to determine recent temperature trends. Please also note that such fits only attempt to describe the past, and usually have small, if any, predictive power. In addition, before using any linear trend (or other) analysis of time series a proper statistical model should be chosen, based on statistical justification.

For temperature time series, there is no *a priori* physical reason why the long-term trend should be linear in time. In fact, climatic time series often have trends for which a straight line is not a good approximation, as can clearly be seen from several of the diagrams in the present report.


For an excellent description of problems often encountered by analyses of temperature time series analyses, please see [Keenan, D.J. 2014: Statistical Analyses of Surface Temperatures in the IPCC Fifth Assessment Report.](#)

**Sunspot activity and QC1 average satellite global air temperature, updated to October 2018**



Variation of global monthly air temperature according to Quality Class 1 (UAH and RSS; see p.4) and observed sunspot number as provided by the Solar Influences Data Analysis Center (SIDC), since 1979. The thin lines represent the monthly values, while the thick line is the simple running 37-month average, nearly corresponding to a running 3-year average. The asymmetrical temperature 'bump' around 1998 is influenced by the oceanographic El Niño phenomenon in 1998, as is the case also for 2015-16.

**2018. The concept of climate revisited.**



*Köppen climate map 2018. Peel, M. C., Finlayson, B. L., and McMahon, T. A. (University of Melbourne) Enhanced, modified, and vectorized by Ali Zifan.*

*“He who controls the past, controls the future. He who controls the present, controls the past” (Orwell 1949).*

The following text is primarily based on and partly extracted from Lamb (1995) and Lindzen (2018). In these two publications various important additional reflections on what constitutes the scientific concept of ‘climate’ can be found.

Climate is the statistics of weather over periods of time (several years), from where good observational

data are available. Thus, climate is always defined from observations from a past time period. The difference between climate and weather is usefully summarized by the popular phrase "Climate is what you expect, weather is what you get."

Climate is measured by assessing the patterns of variation in temperature, humidity, atmospheric

pressure, wind, precipitation, atmospheric particle count and other meteorological variables over long periods of time (several years), averages as well as extremes and frequencies (Lindzen 2018). Climate differs from weather, in that weather only describes the short-term conditions of these variables.

Climate is generated by the climate system, which has five components: atmosphere, hydrosphere, cryosphere, lithosphere, and biosphere (Lindzen 2018). Thus, the climate of a certain location is affected – among other things - by its latitude, terrain, and altitude, as well as nearby oceans. Climates can be classified according to the average and the typical ranges of different variables, most commonly temperature, precipitation and wind. The most widespread climate classification scheme is the Köppen climate classification (see diagram on previous page).

Climate (from Ancient Greek *klima*, meaning *inclination*) is commonly defined as the average weather observed over a long period, as least several years. A common averaging period is 30 years, but other periods may be used depending on the purpose, and the degree of meteorological variability in the region considered. Thus, there is no fixed period for defining climate.

Until around 1960, it was generally assumed that for all practical purposes and decisions climate could be considered constant (Lamb 1995). This view seemed at that time to be soundly based in science. The first long series of regular meteorological observations made in cities of Europe and North America showed that climate of the late 19<sup>th</sup> century to be very similar to the period about a hundred years earlier when the observations had been instituted. Many working practices in applying climatology to forward planning was therefore based on this assumed constancy of climate, even though most geologists and archaeologists at that time realized that climate and environment has a much more interesting history than that (Lamb 1995).

The first formal step in the direction of standardization of the concept 'climate' was taken at the 1935 conference of the International Meteorological Organization (forerunner of the present World Meteorological Organization; WMO) when use of the observations of the years 1901-1930 for all climatic purposes was recommended as the so-called 'climatic normal period'. Use of a 30-year period for defining climate was in this way somewhat accidentally; at that time there just happened to be about 30 years with good meteorological observations available. Also, the choice of the word 'normal' turned out to be rather unfortunate but it has nevertheless persisted in climatological practice.

Regrettably, the term 'climatic normal' to non-experts suggests that nature recognizes such a norm and that conditions should continually return to those of the 'normal period'. This is however a naïve perception of climate, as is today well documented by archaeological, geological, geomorphological and glaciological knowledge on past climates and environments.

Nevertheless, even in 2018, many people still erroneously believe that climate on planet Earth in the absence of humans would essentially be stable and unchanging.

The climate system on planet Earth consists of two turbulent fluids (the atmosphere and oceans) interacting with each other. These fluids are on a rotating planet that is unevenly heated by the sun. The motions in the atmosphere (and to a lesser extent in the oceans) are generated by the uneven influence of the sun. The sun, itself, may display variations, but it always shines directly on the tropics while barely skimming the Earth at the poles. The drivers of the oceans are more complex and include forcing by wind as well as the sinking of cold and salty water. The rotation of the Earth has many consequences, and, among other things, leads to

incoming radiation being distributed around a latitude circle (Lindzen 2018).

The oceans have circulations and currents operating on time scales ranging from years to millennia, and these systems carry heat to and from the surface. The gravity exerted by Sun and Moon are responsible for tidal variations, that also influences on the intensity of certain ocean currents. Because of the scale and density of the oceans, the flow speeds are generally much smaller than in the atmosphere and are associated with much longer timescales. The fact that these circulations carry heat to and from the surface means that the surface, itself, is never in equilibrium with space. There is never an exact balance between incoming heat from the sun and outgoing radiation generated by the Earth because heat is always being stored in and released from the oceans and surface temperature is always, therefore, varying somewhat (Lindzen 2018).

---

51

In addition to the oceans, the atmosphere is interacting with a hugely irregular land surface. As air passes over mountain ranges, the flow is greatly distorted. Topography therefore plays a major role in modifying regional climate. These distorted air-flows even generate fluid waves in the atmosphere that can alter climate at distant locations (Lindzen 2018).

Paleoclimatology is the study of past climates, before observational data are at hand. Since frequent and direct observations of climate are generally not available before the 19th century, paleoclimates are inferred from proxy variables that include non-biotic evidence such as sediments found in lake beds and ice cores, evidence for past glacier activity (e.g. moraines) and permafrost (e.g. ice-wedges and rock glaciers) and biotic evidence such as tree rings and corals. Climate change may occur over long and short timescales, induced by a variety of factors.

Recently, there has been a new and unfortunate development in the understanding of the term 'climate'. For its recent 1.5°C SR15-report the IPCC (IPCC 2018) has changed the definition of climate to what has been loosely called "the climate we are in" (Whitehouse 2018).

Climate is now defined by the IPCC as a speculative 30-year global average temperature that is based, on one hand, on the observed global temperature data from the past 15 years and, on the other hand, on assumed global temperatures for the next 15 years (IPCC 2018, page 6, footnote 5). In other words, this new IPCC definition mixes observed data with speculative data and simply assumes that a short-term 15-year trend won't change for another 15 years in the future (Whitehouse 2018). By this all evidence for periodicity of climate variations playing out over more than 7-8 years are effectively ignored.

This new proposition was put before the recent IPCC meeting at Incheon (Republic of Korea) and agreed as a reasonable thing to do to better communicate climate trends.

Unfortunately, this new IPCC definition of climate is philosophically unsound as it is open to speculation and manipulation, and it therefore remains questionable (Whitehouse 2018). Presumably, if sustained, it is likely to create problems for the IPCC.

The new IPCC definition of climate is mainly a political construct. It is almost certainly good for creating newspaper headlines but has little scientific substance. Extrapolating assumed, non-existing data into the future for calculating temperature trends over the coming 15 years do not make scientific sense and can never be part of a definition of what constitutes climate according to scientific standards. Climate is always defined from observations made in the past.

REFERENCES:

IPCC 2018. Global warming of 1.5°C. SR15. [http://report.ipcc.ch/sr15/pdf/sr15\\_spm\\_final.pdf](http://report.ipcc.ch/sr15/pdf/sr15_spm_final.pdf)

Köppen climate classification 2018. [https://en.wikipedia.org/wiki/Köppen\\_climate\\_classification](https://en.wikipedia.org/wiki/Köppen_climate_classification)

Köppen climate map 2018. Peel, M. C., Finlayson, B. L., and McMahon, T. A. (University of Melbourne) Enhanced, modified, and vectorized by Ali Zifan. - Hydrology and Earth System Sciences: "Updated world map of the Köppen-Geiger climate classification" (Supplement) map in PDF (Institute for Veterinary Public Health). [https://en.wikipedia.org/wiki/Köppen\\_climate\\_classification](https://en.wikipedia.org/wiki/Köppen_climate_classification)

Lindzen, R. 2018. *Global warming for the two cultures*. The GlobalWarming Policy Foundation, 2018 Annual GWPF Lecture, Institution of Mechanical Engineers, London. <https://www.thegwgf.org/richard-lindzen-global-warming-for-the-two-cultures/>

Lamb, H.H. 1995. *Climate history and the modern world*. Second edition, Routledge, London and New York, 433 pp. ISBN 0-415-12735-1.

Orwell, G. 1949. *Nineteen Eighty-Four. A novel*. New York: Harcourt, Brace & Co.

Whitehouse, D. 2018. *Moving the Goalposts, IPCC Secretly Redefines Climate*. GWPF, Oct 29, 2018. <https://www.thegwgf.com/moving-goalposts-ipcc-secretly-redefines-climate/>

Wikipedia 2018. <https://en.wikipedia.org/wiki/Climate>

\*\*\*\*\*

All diagrams in this report, along with any supplementary information, including links to data sources and previous issues of this newsletter, are freely available for download on [www.climate4you.com](http://www.climate4you.com)

Yours sincerely,

Ole Humlum (Ole.Humlum@gmail.com)

Longyearbyen, Svalbard

November 20, 2018.